

# International Journal on Robotics, Automation and Sciences

## Neighbourly Edge Irregularity on Interval-valued Pythagorean Neutrosophic Graph

Murugappan Mullai\*, Govindan Vetrivel, Grienggrai Rajchakit\* and G. Madhan Kumar

**Abstract** – Interval-Valued Pythagorean Neutrosophic Graph (IVPNG) comprises independent and dependent membership elements of interval entries. This study aims to extend the findings of the fuzzy concept to a neutrosophic environment, which is helpful in examining a problem or an event with three associated memberships. This paper demonstrates some kinds of irregular properties on IVPNG, like neighbourly irregular, neighbourly totally irregular, strongly irregular, strongly totally irregular, highly irregular, and highly totally irregular. On certain conditions, neighbourly edge irregular and neighbourly edge totally irregular IVPNG satisfy these irregular properties and vice versa.

**Keywords**— *Neutrosophic Graph, Interval-Valued Pythagorean Neutrosophic Graph, Edge Irregular IVPNG, Neighbourly Edge Irregular IVPNG, Neighbourly Edge Totally Irregular IVPNG.*

### I. INTRODUCTION

Zadeh's contribution [14] of fuzzy set and fuzzy relation led to the foundation of Rosenfeld's fuzzy graph and its applications. NagoorGani and Radha [9] elaborated the properties of fuzzy work by introducing the regularity of fuzzy graphs, which considers the equivalent value in degree and totally degree. The irregularity concept on

fuzzy graphs was initiated and extended to find neighbourly irregular and highly irregular fuzzy graphs by NagoorGani and Latha [10]. Strongly irregular properties on fuzzy graphs and their properties were illustrated in detail by Nandhini and Nandhini [11]. The regularity of edge degree and total edge degree in fuzzy graphs and their extension to interval-valued fuzzy graph (IVFG) was presented by Radha and Kumaravel. Narayanan and Maheswari extended the theory by analyzing the irregularity property on edge, edge totally, strongly edge, and strongly edge totally IVFG. With these sources, Sundarapandian and Ravi Narayanan developed the neighbourly edge irregular and neighborly edge totally irregular IVFG.

Atanassov explored the theory of intuitionistic fuzzy set from a fuzzy set by adding a non-membership function and defined an intuitionistic fuzzy graph. Later, Atanassov and Gargov jointly developed the interval-valued intuitionistic fuzzy set, which was refined into defining the interval-valued intuitionistic fuzzy graph.

Smarandache [12,13] came up with an additional membership function called "indeterminacy" and defined the neutrosophic set, and later, it emerged as a neutrosophic graph. Broumi and Smarandache have worked jointly on Single valued neutrosophic graph and

\*Corresponding author. Email: [mullaim@alagappauniversity.ac.in](mailto:mullaim@alagappauniversity.ac.in), [kreangkri@mju.ac.th](mailto:kreangkri@mju.ac.th), ORCID: 0000-0001-5762-1308  
Murugappan Mullai, Assistant Professor (DDE), Department of Mathematics, Alagappa University, Karaikudi, Tamilnadu, India (e-mail: [mullaim@alagappauniversity.ac.in](mailto:mullaim@alagappauniversity.ac.in)).

Govindan Vetrivel, Department of Mathematics, Alagappa University, Karaikudi, Tamilnadu, India (e-mail: [menakagovindan@gmail.com](mailto:menakagovindan@gmail.com) ).  
Grienggrai Rajchakit, Associate Professor, Department of Mathematics, Faculty of Science, Maejo University, Chiang Mai 50290, Thailand (e-mail: [kreangkri@mju.ac.th](mailto:kreangkri@mju.ac.th)).

G. Madhan Kumar, Department of Mathematics, Alagappa University, Karaikudi, Tamilnadu, India (e-mail: [madhan001kumar@gmail.com](mailto:madhan001kumar@gmail.com)).

International Journal on Robotics, Automation and Sciences (2023) 5,2:54-58

<https://doi.org/10.33093/ijoras.2023.5.2.6>

Manuscript received: 13 May 2023 | Revised: 28 July 2023 | Accepted: 10 August 2023 | Published: 30 September 2023

© Universiti Telekom Sdn Bhd.

Published by MMU PRESS. URL: <http://journals.mmupress.com/ijoras>

This article is licensed under the Creative Commons BY-NC-ND 4.0 International License



PRESS



its properties. Broumi et al. [5,6] coined the term "Interval-valued neutrosophic graph," which is the generalization of the above all mentioned graph types. Ajay and Chellamani [1] restricted the independent memberships of a neutrosophic graph by introducing a Pythagorean neutrosophic fuzzy graph, which bears the dependent component. This paper deals with the initiation of neighbourly edge irregular interval-valued Pythagorean neutrosophic graphs (NEI-IVPNG), neighbourly edge totally irregular interval-valued Pythagorean neutrosophic graphs (NETI-IVPNG), and finally, a comparative study between NEI-IVPNG and NETI-IVPNG is done and also analyzed with SI-IVPNG, HI-IVPNG.

## II. PRELIMINARIES

### Definition 2.1.[1]

A Pythagorean Neutrosophic Graph  $G$  is defined to be a set  $G = (A, B)$ , where

- (i) The functions  $M_A : V \rightarrow [0,1]$ ,  $I_A : V \rightarrow [0,1]$  and  $N_A : V \rightarrow [0,1]$  represent the degree of membership, indeterminacy membership and non-membership of the element  $x \in V$ , respectively, such that  $0 \leq M_A(x)^2 + I_A(x)^2 + N_A(x)^2 \leq 2$ , for all  $x \in V$ ,
- (ii) The functions  $M_B : E \subseteq V \times V \rightarrow [0,1]$ ,  $I_B : E \subseteq V \times V \rightarrow [0,1]$  and  $N_B : E \subseteq V \times V \rightarrow [0,1]$  are defined by  $M_B(xy) \leq \min\{M_A(x), M_A(y)\}$ ,  $I_B(xy) \leq \min\{I_A(x), I_A(y)\}$  and  $N_B(xy) \geq \max\{N_A(x), N_A(y)\}$  such that  $0 \leq M_B(xy)^2 + I_B(xy)^2 + N_B(xy)^2 \leq 2$ , for all  $xy \in E$

### Definition 2.2.

Let  $G = (A, B)$  be an Interval-Valued Pythagorean Neutrosophic Graph (IVPNG), where

- (i) The functions  $M_A : V \rightarrow D[0,1]$ ,  $I_A : V \rightarrow D[0,1]$  and  $N_A : V \rightarrow D[0,1]$  represent the degree of interval-valued membership, interval-valued indeterminacy membership and interval-valued non-membership of the element  $x \in V$ , respectively, such that  $M_{A+}(x)^2 + I_{A+}(x)^2 + N_{A+}(x)^2 \leq 2$ ,  $M_{A-}(x)^2 + I_{A-}(x)^2 + N_{A-}(x)^2 \leq 2$  for all  $x \in V$ ,
- (ii) The functions  $M_B : E \subseteq V \times V \rightarrow D[0,1]$ ,  $I_B : E \subseteq V \times V \rightarrow D[0,1]$  and  $N_B : E \subseteq V \times V \rightarrow D[0,1]$  are defined by  $M_{B-}(xy) \leq \min\{M_{A-}(x), M_{A-}(y)\}$ ;  $M_{B+}(xy) \leq \min\{M_{A+}(x), M_{A+}(y)\}$ ;  $I_{B-}(xy) \leq \min\{I_{A-}(x), I_{A-}(y)\}$ ;  $I_{B+}(xy) \leq \min\{I_{A+}(x), I_{A+}(y)\}$ ;  $N_{B-}(xy) \geq \max\{N_{A-}(x), N_{A-}(y)\}$ ;  $N_{B+}(xy) \geq \max\{N_{A+}(x), N_{A+}(y)\}$  such that  $M_{B+}(xy)^2 + I_{B+}(xy)^2 + N_{B+}(xy)^2 \leq 2$ ,  $M_{B-}(xy)^2 + I_{B-}(xy)^2 + N_{B-}(xy)^2 \leq 2$  for all  $xy \in E$ . Here,  $A$  is the interval-valued pythagorean neutrosophic vertex set of  $G$  and  $B$  is the interval-valued pythagorean neutrosophic edge set of  $G$ .

### Defintion 2.3.

If all set of adjacent edges of a connected IVPNG have distinct degree then the graph is known as NEI-IVPNG.

### Definition 2.4.

If all set of adjacent edges of a connected IVPNG have distinct total degree then the graph is known as NETI-IVPNG.

### Example 2.5.

Let  $G: (V, E)$  be an IVPNG. From Figure 1,  $d(u) = d(v) = d(w) = d(x) = \langle (0.3, 0.5), (0.7, 0.7), (0.5, 1) \rangle$ ,  $d(uv) = d(wx) = \langle (0.4, 0.6), (0.8, 0.8), (0.4, 0.8) \rangle$  and  $d(vw) = d(xu) = \langle (0.2, 0.4), (0.6, 0.6), (0.6, 1.2) \rangle$ .

It shows that, all set of adjacent edges have distinct degree.

Therefore,  $G$  is concluded as an NEI-IVPNG.

Total degrees of the edges are listed as follows:

$td(uv) = td(wx) = \langle (0.5, 0.8), (1.1, 1.1), (0.7, 1.4) \rangle$ ,

$td(vw) = td(xu) = \langle (0.4, 0.7), (1.0, 1.0), (0.8, 1.6) \rangle$ .

It shows that, all set of adjacent edges have distinct total degree. Therefore,  $G$  is a NETI-IVPNG.

Hence, the graph  $G$  is both NEI-IVPNG and NETI-IVPNG.

The following graph is an example for both NEI-IVPNG and NETI-IVPN:




FIGURE 1. Graph for NEI-IVPNG and NETI-IVPN.

### Definition 2.6.

Let  $G = (A, B)$  denotes a connected IVPNG. Then  $G$  is said to be a strongly irregular interval valued pythagorean neutrosophic graph (SI-IVPNG) if all set of vertices in  $G$  have distinct degrees (or) no two vertices have same degree.

### Definition 2.7.

Let  $G = (A, B)$  denotes a connected IVPNG. Then  $G$  is highly irregular interval-valued pythagorean neutrosophic graph (HI-IVPNG) if all vertex is adjacent to the vertices in  $G$  have distinct degree.

## III. MAIN RESULTS

### Theorem 3.1.

Let  $G$  be a connected IVNPG and  $B$  be a constant function.  $G$  is a NEI-IVPNG if and only if  $G$  is a NETI-IVPNG.

### Proof:

Consider a constant function  $B$ . Let  $B(uv) = \langle (M_{B-}(uv), M_{B+}(uv)), (I_{B-}(uv), I_{B+}(uv)), (N_{B-}(uv), N_{B+}(uv)) \rangle = \langle (c_1, c_2), (b_1, b_2), (d_1, d_2) \rangle$ ,  $\forall uv \in E$ , where  $c_1, c_2, b_1, b_2, d_1, d_2$  are constants.

Let  $G$  be a NEI-IVPNG.

Then, the set of all adjacent edges has distinct degrees.

$\Leftrightarrow d(uv) \neq d(vw)$ , where  $E = \{uv, vw\}$  are adjacent edges.

$\Leftrightarrow \langle d_M(uv), d_I(uv), d_N(uv) \rangle \neq \langle d_M(vw), d_I(vw), d_N(vw) \rangle$   
 $\Leftrightarrow d_M(uv) \neq d_M(vw)$  (or)  $d_I(uv) \neq d_I(vw)$  (or)  $d_N(uv) \neq d_N(vw)$  (or) all of them.

Without loss of generality, let us assume that  $d_M(uv) \neq d_M(vw)$

$\Leftrightarrow (d_{M-}(uv), d_{M+}(uv)) \neq (d_{M-}(vw), d_{M+}(vw))$

$\Leftrightarrow d_{M-}(uv) \neq d_{M-}(vw)$  (or)  $d_{M+}(uv) \neq d_{M+}(vw)$  (or) both.

Without loss of generality, let us assume that  $d_{M-}(uv) \neq d_{M-}(vw)$

$\Leftrightarrow d_{M-}(u) + d_{M-}(v) - 2M_{B-}(uv) \neq d_{M-}(v) + d_{M-}(w) - 2M_{B-}(vw)$

$\Leftrightarrow d_{M-}(u) + d_{M-}(v) - 2c_1 \neq d_{M-}(v) + d_{M-}(w) - 2c_1$

Adding  $c_1$  on both sides,

$\Leftrightarrow d_{M-}(u) + d_{M-}(v) - 2c_1 + c_1 \neq d_{M-}(v) + d_{M-}(w) - 2c_1 + c_1$

$\Leftrightarrow d_{M-}(u) + d_{M-}(v) - c_1 \neq d_{M-}(v) + d_{M-}(w) - c_1$

$\Leftrightarrow d_{M-}(u) + d_{M-}(v) - M_{B-}(uv) \neq d_{M-}(v) + d_{M-}(w) - M_{B-}(vw)$

$\Leftrightarrow td_{M-}(uv) \neq td_{M-}(vw) \Leftrightarrow td_M(uv) \neq td_M(vw)$

$\Leftrightarrow \langle td_M(uv), td_I(uv), td_N(uv) \rangle \neq \langle td_M(vw), td_I(vw), td_N(vw) \rangle$

$\Leftrightarrow td(uv) \neq td(vw)$ , where  $E = \{uv, vw\}$  are adjacent edges.

Then, the set of all adjacent edges has distinct total degrees.

$\Leftrightarrow$  Hence,  $G$  is a NETI-IVPNG.

### Remark 3.2.

Let  $G$  be a connected IVPNG. If  $G$  is both NEI-IVPNG and NETI-IVPNG, then  $B$  need not to be a constant function.

### Theorem 3.3.

Let  $G$  be a connected IVPNG and  $B$  is a constant function. If  $G$  is a SI-IVPNG, then  $G$  is a NEI-IVPNG.

#### Proof:

Let  $B(uv) = \langle (M_{B-}(uv), M_{B+}(uv)), (I_{B-}(uv), I_{B+}(uv)), (N_{B-}(uv), N_{B+}(uv)) \rangle = \langle (c_1, c_2), (b_1, b_2), (d_1, d_2) \rangle$ ,  $\forall uv \in E$ , where  $c_1, c_2, b_1, b_2, d_1, d_2$  are constants. Let  $uv$  and  $vw$  are any two adjacent edges in  $G$ . Assume that  $G$  is a SI-IVPNG. Then, all pair of vertices in  $G$  having distinct degrees.

$\Rightarrow d(u) \neq d(v) \neq d(w) \Rightarrow d(u) \neq d(w)$

Adding  $d(v)$  on both sides,

$\Rightarrow d(u) + d(v) \neq d(v) + d(w)$

$\Rightarrow \langle (d_{M-}(u), d_{M+}(u)), (d_{I-}(u), d_{I+}(u)), (d_{N-}(u), d_{N+}(u)) \rangle + \langle (d_{M-}(v), d_{M+}(v)), (d_{I-}(v), d_{I+}(v)), (d_{N-}(v), d_{N+}(v)) \rangle \neq \langle (d_{M-}(v), d_{M+}(v)), (d_{I-}(v), d_{I+}(v)), (d_{N-}(v), d_{N+}(v)) \rangle + \langle (d_{M-}(w), d_{M+}(w)), (d_{I-}(w), d_{I+}(w)), (d_{N-}(w), d_{N+}(w)) \rangle$

Adding  $-2\langle (c_1, c_2), (b_1, b_2), (d_1, d_2) \rangle$  on both sides,  
 $\Rightarrow \langle (d_{M-}(u) + d_{M-}(v) - 2c_1, d_{M+}(u) + d_{M+}(v) - 2c_2), (d_{I-}(u) + d_{I-}(v) - 2b_1, d_{I+}(u) + d_{I+}(v) - 2b_2), (d_{N-}(u) + d_{N-}(v) - 2d_1, d_{N+}(u) + d_{N+}(v) - 2d_2) \rangle \neq \langle (d_{M-}(v) + d_{M-}(w) - 2c_1, d_{M+}(v) + d_{M+}(w) - 2c_2), (d_{I-}(v) + d_{I-}(w) - 2b_1, d_{I+}(v) + d_{I+}(w) - 2b_2), (d_{N-}(v) + d_{N-}(w) - 2d_1, d_{N+}(v) + d_{N+}(w) - 2d_2) \rangle$

$\Rightarrow \langle (d_{M-}(u) + d_{M-}(v) - 2M_{B-}(uv), d_{M+}(u) + d_{M+}(v) - 2M_{B+}(uv)), (d_{I-}(u) + d_{I-}(v) - 2I_{B-}(uv), d_{I+}(u) + d_{I+}(v) - 2I_{B+}(uv)), (d_{N-}(u) + d_{N-}(v) - 2N_{B-}(uv), d_{N+}(u) + d_{N+}(v) - 2N_{B+}(uv)) \rangle \neq \langle (d_{M-}(v) + d_{M-}(w) - 2M_{B-}(vw), d_{M+}(v) + d_{M+}(w) - 2M_{B+}(vw)), (d_{I-}(v) + d_{I-}(w) - 2I_{B-}(vw), d_{I+}(v) + d_{I+}(w) - 2I_{B+}(vw)), (d_{N-}(v) + d_{N-}(w) - 2N_{B-}(vw), d_{N+}(v) + d_{N+}(w) - 2N_{B+}(vw)) \rangle$   
 $\Rightarrow \langle (d_{M-}(uv), d_{M+}(uv)), (d_{I-}(uv), d_{I+}(uv)), (d_{N-}(uv), d_{N+}(uv)) \rangle \neq \langle (d_{M-}(vw), d_{M+}(vw)), (d_{I-}(vw), d_{I+}(vw)), (d_{N-}(vw), d_{N+}(vw)) \rangle$   
 $\Rightarrow d(uv) \neq d(vw)$

Hence, the set of all adjacent edges has distinct degree.

This implies that,  $G$  is a NEI-IVPNG.

### Theorem 3.4.

Let  $G$  be a connected IVPNG and  $B$  is a constant function. If  $G$  is a SI-IVPNG, then  $G$  is a NETI-IVPNG.

#### Proof:

Let  $G$  be a connected IVPNG and  $B$  be a constant function. Suppose that  $G$  is SI-IVPNG. By the above theorem,  $G$  is a NEI-IVPNG. From theorem 3.1,  $G$  is a NETI-IVPNG.

### Remark 3.5.

The converse part of the theorem 3.3 and 3.4 need not be true.

### Theorem 3.6.

Let  $G$  be a connected IVPNG and  $B$  denotes a constant function. Then,  $G$  is a HI-IVPNG iff  $G$  is a NEI-IVPNG.

**Proof:** Let  $B(uv) = \langle (M_{B-}(uv), M_{B+}(uv)), (I_{B-}(uv), I_{B+}(uv)), (N_{B-}(uv), N_{B+}(uv)) \rangle = \langle (c_1, c_2), (b_1, b_2), (d_1, d_2) \rangle$ ,  $\forall uv \in E$ , where  $c_1, c_2, b_1, b_2, d_1, d_2$  are constants.

Consider the adjacent edges  $uv$  and  $vw$  in  $G$  and let  $G$  be a HI-IVPNG. Then, by definition 2.7,

$\Rightarrow d(u) \neq d(w)$

Adding  $d(v)$  on both sides,

$\Rightarrow d(u) + d(v) \neq d(v) + d(w)$

$\Rightarrow \langle (d_{M-}(u), d_{M+}(u)), (d_{I-}(u), d_{I+}(u)), (d_{N-}(u), d_{N+}(u)) \rangle + \langle (d_{M-}(v), d_{M+}(v)), (d_{I-}(v), d_{I+}(v)), (d_{N-}(v), d_{N+}(v)) \rangle \neq \langle (d_{M-}(v), d_{M+}(v)), (d_{I-}(v), d_{I+}(v)), (d_{N-}(v), d_{N+}(v)) \rangle + \langle (d_{M-}(w), d_{M+}(w)), (d_{I-}(w), d_{I+}(w)), (d_{N-}(w), d_{N+}(w)) \rangle$

Adding  $-2\langle (c_1, c_2), (b_1, b_2), (d_1, d_2) \rangle$  on both sides,

$$\begin{aligned}
&\Rightarrow \langle (d_{M-}(u) + d_{M-}(v) - 2c_1, d_{M+}(u) + d_{M+}(v) - 2c_2), (d_{I-}(u) + d_{I-}(v) - 2b_1, d_{I+}(u) + d_{I+}(v) - 2b_2), (d_{N-}(u) + d_{N-}(v) - 2d_1, d_{N+}(u) + d_{N+}(v) - 2d_2) \rangle \neq \langle (d_{M-}(v) + d_{M-}(w) - 2c_1, d_{M+}(v) + d_{M+}(w) - 2c_2), (d_{I-}(v) + d_{I-}(w) - 2b_1, d_{I+}(v) + d_{I+}(w) - 2b_2), (d_{N-}(v) + d_{N-}(w) - 2d_1, d_{N+}(v) + d_{N+}(w) - 2d_2) \rangle \\
&\Rightarrow \langle (d_{M-}(u) + d_{M-}(v) - 2M_{B-}(uv), d_{M+}(u) + d_{M+}(v) - 2M_{B+}(uv)), (d_{I-}(u) + d_{I-}(v) - 2I_{B-}(uv), d_{I+}(u) + d_{I+}(v) - 2I_{B+}(uv)), (d_{N-}(u) + d_{N-}(v) - 2N_{B-}(uv), d_{N+}(u) + d_{N+}(v) - 2N_{B+}(uv)) \rangle \\
&\neq \langle (d_{M-}(v) + d_{M-}(w) - 2M_{B-}(vw), d_{M+}(v) + d_{M+}(w) - 2M_{B+}(vw)), (d_{I-}(v) + d_{I-}(w) - 2I_{B-}(vw), d_{I+}(v) + d_{I+}(w) - 2I_{B+}(vw)), (d_{N-}(v) + d_{N-}(w) - 2N_{B-}(vw), d_{N+}(v) + d_{N+}(w) - 2N_{B+}(vw)) \rangle \\
&\Rightarrow \langle (d_{M-}(uv), d_{M+}(uv)), (d_{I-}(uv), d_{I+}(uv)), (d_{N-}(uv), d_{N+}(uv)) \rangle \neq \langle (d_{M-}(vw), d_{M+}(vw)), (d_{I-}(vw), d_{I+}(vw)), (d_{N-}(vw), d_{N+}(vw)) \rangle \\
&\Rightarrow d(uv) \neq d(vw)
\end{aligned}$$

Hence, all set of adjacent edges have distinct degree. Therefore, G is a NEI-IVPNG.

Conversely, assume that G is a NEI-IVPNG. All set of adjacent edges in G has distinct degrees. Let v be the vertex which is adjacent to u, w and x. Then we have uv, vw and vx as the adjacent edges in G.

$$\begin{aligned}
&\Rightarrow d(uv) \neq d(vw) \neq d(vx) \\
&\Rightarrow \langle (d_{M-}(uv), d_{M+}(uv)), (d_{I-}(uv), d_{I+}(uv)), (d_{N-}(uv), d_{N+}(uv)) \rangle \neq \langle (d_{M-}(vw), d_{M+}(vw)), (d_{I-}(vw), d_{I+}(vw)), (d_{N-}(vw), d_{N+}(vw)) \rangle \neq \langle (d_{M-}(vx), d_{M+}(vx)), (d_{I-}(vx), d_{I+}(vx)), (d_{N-}(vx), d_{N+}(vx)) \rangle \\
&\Rightarrow \langle (d_{M-}(u) + d_{M-}(v) - 2M_{B-}(uv), d_{M+}(u) + d_{M+}(v) - 2M_{B+}(uv)), (d_{I-}(u) + d_{I-}(v) - 2I_{B-}(uv), d_{I+}(u) + d_{I+}(v) - 2I_{B+}(uv)), (d_{N-}(u) + d_{N-}(v) - 2N_{B-}(uv), d_{N+}(u) + d_{N+}(v) - 2N_{B+}(uv)) \rangle \neq \langle (d_{M-}(v) + d_{M-}(w) - 2M_{B-}(vw), d_{M+}(v) + d_{M+}(w) - 2M_{B+}(vw)), (d_{I-}(v) + d_{I-}(w) - 2I_{B-}(vw), d_{I+}(v) + d_{I+}(w) - 2I_{B+}(vw)), (d_{N-}(v) + d_{N-}(w) - 2N_{B-}(vw), d_{N+}(v) + d_{N+}(w) - 2N_{B+}(vw)) \rangle \neq \langle (d_{M-}(v) + d_{M-}(x) - 2M_{B-}(vx), d_{M+}(v) + d_{M+}(x) - 2M_{B+}(vx)), (d_{I-}(v) + d_{I-}(x) - 2I_{B-}(vx), d_{I+}(v) + d_{I+}(x) - 2I_{B+}(vx)), (d_{N-}(v) + d_{N-}(x) - 2N_{B-}(vx), d_{N+}(v) + d_{N+}(x) - 2N_{B+}(vx)) \rangle
\end{aligned}$$

$$\begin{aligned}
&\Rightarrow \langle (d_{M-}(u) + d_{M-}(v) - 2c_1, d_{M+}(u) + d_{M+}(v) - 2c_2), (d_{I-}(u) + d_{I-}(v) - 2b_1, d_{I+}(u) + d_{I+}(v) - 2b_2), (d_{N-}(u) + d_{N-}(v) - 2d_1, d_{N+}(u) + d_{N+}(v) - 2d_2) \rangle \neq \langle (d_{M-}(v) + d_{M-}(w) - 2c_1, d_{M+}(v) + d_{M+}(w) - 2c_2), (d_{I-}(v) + d_{I-}(w) - 2b_1, d_{I+}(v) + d_{I+}(w) - 2b_2), (d_{N-}(v) + d_{N-}(w) - 2d_1, d_{N+}(v) + d_{N+}(w) - 2d_2) \rangle \neq \langle (d_{M-}(v) + d_{M-}(x) - 2c_1, d_{M+}(v) + d_{M+}(x) - 2c_2), (d_{I-}(v) + d_{I-}(x) - 2b_1, d_{I+}(v) + d_{I+}(x) - 2b_2), (d_{N-}(v) + d_{N-}(x) - 2d_1, d_{N+}(v) + d_{N+}(x) - 2d_2) \rangle
\end{aligned}$$

Adding  $2\langle (c_1, c_2), (b_1, b_2), (d_1, d_2) \rangle$  on both sides,

$$\begin{aligned}
&\Rightarrow \langle (d_{M-}(u), d_{M+}(u)), (d_{I-}(u), d_{I+}(u)), (d_{N-}(u), d_{N+}(u)) \rangle + \langle (d_{M-}(v), d_{M+}(v)), (d_{I-}(v), d_{I+}(v)), (d_{N-}(v), d_{N+}(v)) \rangle \neq \langle (d_{M-}(v), d_{M+}(v)), (d_{I-}(v), d_{I+}(v)), (d_{N-}(v), d_{N+}(v)) \rangle + \langle (d_{M-}(w), d_{M+}(w)), (d_{I-}(w), d_{I+}(w)), (d_{N-}(w), d_{N+}(w)) \rangle \neq \langle (d_{M-}(v), d_{M+}(v)), (d_{I-}(v), d_{I+}(v)), (d_{N-}(v), d_{N+}(v)) \rangle + \langle (d_{M-}(x), d_{M+}(x)), (d_{I-}(x), d_{I+}(x)), (d_{N-}(x), d_{N+}(x)) \rangle \\
&\Rightarrow d(u) + d(v) \neq d(v) + d(w) \neq d(v) + d(x)
\end{aligned}$$

$$\Rightarrow d(u) \neq d(w) \neq d(x).$$

$\Rightarrow$  Every vertex in G is adjacent to the vertices in G, which have distinct degrees. Hence, G is a HI-IVPNG.

### Theorem 3.7.

Let G be a connected IVPNG and B denotes a constant function. Then, G is an HI-IVPNG, iff G is a NETI-IVPNG.

### Proof:

Let G be a connected IVPNG. Assume that B is a constant function and G is HI-IVPNG.

$\Leftrightarrow$  G is a NEI-IVPNG (from above theorem).

$\Leftrightarrow$  G is a NETI-IVPNG (from Theorem 3.1).

### Theorem 3.8.

Let G be a connected IVPNG and B denotes a constant function. If G is a NI-IVPNG, then G is a NEI-IVPNG.

### Proof:

Let G be a connected IVPNG and B denotes a constant function. Let  $B(uv) = \langle (M_{B-}(uv), M_{B+}(uv)), (I_{B-}(uv), I_{B+}(uv)), (N_{B-}(uv), N_{B+}(uv)) \rangle = \langle (c_1, c_2), (b_1, b_2), (d_1, d_2) \rangle$ ,  $\forall uv \in E$ , where  $c_1, c_2, b_1, b_2, d_1, d_2$  are constants. Consider any two adjacent edges uv and vw in G. Suppose that G is NI-IVPNG.

$\Rightarrow$  The set of all adjacent vertices in G has distinct degree.

$$\Rightarrow d(u) \neq d(v) \text{ and } d(v) \neq d(w)$$

$$\Rightarrow d(u) \neq d(w)$$

Adding  $d(v)$  on both sides,

$$\Rightarrow d(u) + d(v) \neq d(v) + d(w)$$

$$\begin{aligned}
&\Rightarrow \langle (d_{M-}(u), d_{M+}(u)), (d_{I-}(u), d_{I+}(u)), (d_{N-}(u), d_{N+}(u)) \rangle + \langle (d_{M-}(v), d_{M+}(v)), (d_{I-}(v), d_{I+}(v)), (d_{N-}(v), d_{N+}(v)) \rangle \neq \langle (d_{M-}(v), d_{M+}(v)), (d_{I-}(v), d_{I+}(v)), (d_{N-}(v), d_{N+}(v)) \rangle + \langle (d_{M-}(w), d_{M+}(w)), (d_{I-}(w), d_{I+}(w)), (d_{N-}(w), d_{N+}(w)) \rangle
\end{aligned}$$

Subtracting  $2\langle (c_1, c_2), (b_1, b_2), (d_1, d_2) \rangle$  on both sides,

$$\begin{aligned}
&\Rightarrow \langle (d_{M-}(u) + d_{M-}(v) - 2c_1, d_{M+}(u) + d_{M+}(v) - 2c_2), (d_{I-}(u) + d_{I-}(v) - 2b_1, d_{I+}(u) + d_{I+}(v) - 2b_2), (d_{N-}(u) + d_{N-}(v) - 2d_1, d_{N+}(u) + d_{N+}(v) - 2d_2) \rangle \neq \langle (d_{M-}(v) + d_{M-}(w) - 2c_1, d_{M+}(v) + d_{M+}(w) - 2c_2), (d_{I-}(v) + d_{I-}(w) - 2b_1, d_{I+}(v) + d_{I+}(w) - 2b_2), (d_{N-}(v) + d_{N-}(w) - 2d_1, d_{N+}(v) + d_{N+}(w) - 2d_2) \rangle
\end{aligned}$$

$$\begin{aligned}
&\Rightarrow \langle (d_{M-}(u) + d_{M-}(v) - 2M_{B-}(uv), d_{M+}(u) + d_{M+}(v) - 2M_{B+}(uv)), (d_{I-}(u) + d_{I-}(v) - 2I_{B-}(uv), d_{I+}(u) + d_{I+}(v) - 2I_{B+}(uv)), (d_{N-}(u) + d_{N-}(v) - 2N_{B-}(uv), d_{N+}(u) + d_{N+}(v) - 2N_{B+}(uv)) \rangle \neq \langle (d_{M-}(v) + d_{M-}(w) - 2M_{B-}(vw), d_{M+}(v) + d_{M+}(w) - 2M_{B+}(vw)), (d_{I-}(v) + d_{I-}(w) - 2I_{B-}(vw), d_{I+}(v) + d_{I+}(w) - 2I_{B+}(vw)), (d_{N-}(v) + d_{N-}(w) - 2N_{B-}(vw), d_{N+}(v) + d_{N+}(w) - 2N_{B+}(vw)) \rangle
\end{aligned}$$

$$\begin{aligned}
&\Rightarrow \langle (d_{M-}(uv), d_{M+}(uv)), (d_{I-}(uv), d_{I+}(uv)), (d_{N-}(uv), d_{N+}(uv)) \rangle \neq \langle (d_{M-}(vw), d_{M+}(vw)), (d_{I-}(vw), d_{I+}(vw)), (d_{N-}(vw), d_{N+}(vw)) \rangle
\end{aligned}$$

All set of adjacent edges in  $G$  has distinct degrees.  
 Therefore,  $G$  is a NEI-IVPNG.

### Theorem 3.9.

Let  $G$  be a connected IVPNG and  $B$  is a constant function. If  $G$  is a NI-IVPNG, then  $G$  is a NETI-IVPNG.

### Proof:

Let  $G$  be a connected IVPNG. Assume that  $B$  is a constant function. Suppose that  $G$  is NI-IVPNG.  
 $\Leftrightarrow G$  is NEI-IVPNG (from the above theorem).  
 $\Leftrightarrow G$  is NETI-IVPNG (from Theorem 3.1).

### Remark 3.10.

The converse part of the theorem 3.8 and 3.9 need not be true.

## IV. CONCLUSION

In this article, we explored the neighbourly edge and neighbourly edge totally irregularities on interval valued pythagorean neutrosophic graph and the relation with other irregular property-bearing graphs. Though there are some real-life applications related to IVPNG [7] but neighbourly edge irregularity was not applied till now. In the future, we plan to execute further development with applications in this area.

## ACKNOWLEDGMENT

We thank the anonymous reviewers for the careful review of our manuscript.

## FUNDING STATEMENT

The article has been written with the joint financial support of RUSA-Phase 2.0 grant sanctioned vide letter No.F.24-51/2014-U, Policy (TN Multi-Gen), Dept. of Edn.Govt. of India, Dt. 09.10.2018, UGC-SAP (DRS-I) vide letter No.F.510/8/DRS-I/2016(SAP-I) Dt. 23.08.2016 and DST (FST-level I) 657876570 vide letter No.SR/FIST/MS-I/2018/17 Dt. 20.12.2018.

## AUTHOR CONTRIBUTIONS

Murugappan Mullai: Conceptualization, Validation, Writing, Administration;

Govindan Vetrivel: Conceptualization, Validation, Writing, Administration;

Grienggrai Rajchakit: Conceptualization, Validation, Writing, Administration;

G. Madhan Kumar: Conceptualization, Validation, Writing, Administration.

## CONFLICT OF INTERESTS

No conflict of interests were disclosed.

Our publication ethics follow The Committee of Publication Ethics (COPE) guideline.  
<https://publicationethics.org/>

## REFERENCES

- [1] D. Ajay and P. Chellamani, "Pythagorean Neutrosophic Fuzzy Graphs," International Journal of Neutrosophic Science, vol. 11, pp. 108–114, Mar. 2020.  
DOI: <https://doi.org/10.5281/zenodo.4172124>.
- [2] D. Ajay, P. Chellamani, G. Rajchakit, N. Boonsatit, and P. Hammachukiattikul, "Regularity of Pythagorean Neutrosophic Fuzzy Graphs with an Illustration in MCDM," AIMS Mathematics, vol. 7, no. 5, pp. 9424–9442, Mar. 2022.  
DOI: <https://doi.org/10.3934/math.2022523>.
- [3] M. Akram and W. A. Dudek, "Interval-valued Fuzzy Graph," Computer and Mathematics with Applications, vol. 61, no. 2, pp. 289–299, Jan. 2011.  
DOI: <https://doi.org/10.1016/j.camwa.2010.11.004>.
- [4] M. Akram, S. Naz, and B. Davvaz, "Simplified Interval-valued Pythagorean Fuzzy Graphs with Application," Complex & Intelligent Systems, vol. 5, pp. 229–253, May 2019.  
DOI: <https://doi.org/10.1007/s40747-019-0106-3>.
- [5] S. Broumi, M. Talea, A. Bakali, and F. Smarandache, "Interval Valued Neutrosophic Graphs," Critical Review, XII, pp. 5–33, 2016.  
URL: <https://fs.unm.edu/IntervalValuedNeutrosophicGraphs-CR12.pdf> (accessed: 25 July 2023)..
- [6] S. Broumi, F. Smarandache, M. Talea, and A. Bakali, "Operations on Interval Valued Neutrosophic Graphs," in New Trends in Neutrosophic Theory and Applications, edited by F. Smarandache and S. Pramanik, pp. 231–254, 2016.  
URL: [https://www.researchgate.net/publication/311545273\\_Operations\\_on\\_Interval\\_Valued\\_Neutrosophic\\_Graphs](https://www.researchgate.net/publication/311545273_Operations_on_Interval_Valued_Neutrosophic_Graphs) (Accessed: 20 July 2023).
- [7] S. Broumi, R. Sundareswaran, M. Shanmugapriya, P. Chellamani, A. Bakali, and M. Talea, "Determination of Various Factors to Evaluate a Successful Curriculum Design Using Interval-Valued Pythagorean Neutrosophic Graphs," Research Square (preprint), Mar. 2023.  
DOI: <https://doi.org/10.1007/s00500-023-08996-y>.
- [8] J. Mohamed and S. Rajamanickam, "Strong Interval-Valued Pythagorean Fuzzy Soft Graphs," Ratio Mathematica, [S.I.], vol. 46, Mar. 2023.  
DOI: <https://doi.org/10.23755/rm.v46i0.1067>.
- [9] A. Nagoorgani and K. Radha, "On Regular Fuzzy Graphs," Journal of Physical Sciences, vol. 12, pp. 33–40, 2008.  
URL: [https://www.researchgate.net/publication/254399182\\_On\\_Regular\\_Fuzzy\\_Graphs](https://www.researchgate.net/publication/254399182_On_Regular_Fuzzy_Graphs) (Accessed: 5 July, 2023)
- [10] A. Nagoor Gani and S. R. Latha, "On Irregular Fuzzy Graphs," Applied Mathematical Sciences, vol. 6, pp. 517–523, 2012.  
DOI: <https://doi.org/10.13140/2.1.2226.2409>.
- [11] S. P. Nandhini and E. Nandhini, "Strongly Irregular Fuzzy Graphs," International Journal of Mathematical Archive, vol. 5, no. 5, pp. 110–114, 2014.  
URL: <https://acadpubl.eu/iji/2017-112-5/9/9.pdf> (Accessed: 10 July, 2023)
- [12] F. Smarandache, "Neutrosophic Set – A Generalization of the Intuitionistic Fuzzy Set," in Granular Computing, IEEE International Conference, pp. 38–42, 2006.  
DOI: <https://doi.org/10.1109/GRC.2006.1635754>.
- [13] F. Smarandache, "A Geometric Interpretation of the Neutrosophic Set – A Generalization of the Intuitionistic Fuzzy Set," in Granular Computing, IEEE International Conference, pp. 602–606, 2011.  
DOI: <https://doi.org/10.1109/GRC.2011.6122665>.
- [14] L. A. Zadeh, "Fuzzy Sets," Information and Control, vol. 8, no. 3, pp. 338–353, 1965.  
DOI: [https://doi.org/10.1016/S0019-9958\(65\)90241-X](https://doi.org/10.1016/S0019-9958(65)90241-X).